



# Investigating the Effect of Hydroalcoholic Extract of Parsley Leaves (*Petroselinum crispum*) on Anxiety in Rats Treated with Lead Acetate

Fatemeh Bastampoor<sup>1</sup>, Seyed Ebrahim Hosseini<sup>1,2\*</sup>, Mehrdad Shariati<sup>1</sup>, Mokhtar Mokhtari<sup>1</sup>

<sup>1</sup>Department of Biology, Kazerun Branch, Islamic Azad University, Kazerun, Iran

<sup>2</sup>Department of Biology, Faculty of Sciences, Zand Institute of Higher Education, Shiraz, Iran

Received: 17/Jan/2023

Revised: 20/Feb/2023

Accepted: 01/Mar/2023

## Abstract

**Background and aim:** Heavy metals such as lead can have negative effects on the nervous system. Anxiety is one of the most common neurological disorders that can be induced under the influence of lead. The purpose of this study was to investigate the protective effect of the hydroalcoholic extract of parsley leaves (*Petroselinum crispum*; HEPL) on anxiety in rats treated with lead acetate.

**Materials and Methods:** Male rats were divided into 6 groups of 10. The control group did not receive drug treatment, but the sham group animals received 1 ml of distilled water as lead acetate solvent by intraperitoneal injection. The animals of the Lead group received lead acetate at a dose of 20 mg/kg and the animals of the HEPL200 group received HEPL at a dose of 200 mg/kg by intraperitoneal injection. The animals of the Lead+HEPL100 and Lead+HEPL200 groups received lead acetate with a dose of 20 mg/kg as an intraperitoneal injection 2 hours before the intraperitoneal injection of HEPL with a dose of 100 and 200 mg/kg, respectively. The study period in all groups was 21 days. At the end of the study, the open field test was used to measure anxiety behavior including the parameters of Grooming (cleaning the body, face, and legs), Rearing (standing on two rear legs), and Crossing (crossing the lines). Statistical analysis was performed using a one-way analysis of variance and the Tukey post hoc test.

**Results:** The results of this study showed that there was a significant decrease in the number of Grooming, Rearing, and Crossing in the Lead group compared to the control and control groups ( $P < 0.05$ ). In the Lead+HEPL200 group, a significant increase in the number of Grooming, Rearing, and Crossing was observed compared to the Lead group ( $P < 0.05$ ).

**Conclusion:** Lead can cause anxiety behaviors in male rats. However, the administration of HEPL can improve anxiety behaviors in male rats treated with lead acetate.

**Keywords:** Lead, Hydroalcoholic extract, Parsley, Anxiety, Rat

**Cite this article as:** Fatemeh Bastampoor, Seyed Ebrahim Hosseini, Mehrdad Shariati, Mokhtar Mokhtari. Investigating the effect of hydroalcoholic extract of parsley leaves (*Petroselinum crispum*) on anxiety in rats treated with lead acetate. J Altern Vet Med. 2023; 6(16): 929-938.

## \* Corresponding Author

Department of Biology, Faculty of Sciences, Zand Institute of Higher Education, Shiraz, Iran.

E-mail: [ebrahim.hossini@yahoo.com](mailto:ebrahim.hossini@yahoo.com), Orcid: <https://orcid.org/0000-0001-5347-1824>



## Introduction

Neurological diseases and disorders are rarely unidimensional or unifactorial. Even those diseases whose etiology appears to be closely related to genetic factors are usually the product of multiple and intertwined risk factors, of which exposure to environmental chemicals may be a component (Weiss *et al.*, 2008). The public health burden of psychiatric disorders such as depression and anxiety is high, with approximately 450 million people worldwide suffering from psychiatric disorders. The term anxiety describes the experience of worry, apprehension, or nervousness associated with physical, cognitive, and behavioral symptoms. Anxiety may occasionally be experienced as part of normal life and maybe adaptive if it increases readiness for new situations. If the anxiety symptoms are persistent, excessive, or dysfunctional, it can cause neurological and psychological disorders (Gale & Millichamp, 2011). Neurotransmitters play an important role in anxiety. One of the most important neurotransmitters is gabargic neurons, which are responsible for anxiety phenomena. Oxidative stress can also be related to anxiety (Ghasemzadeh Dehkordi *et al.*, 2022). Exposure to heavy metals such as lead during pregnancy and infancy causes widespread neurological and psychological effects such as depression, anxiety, mood disorders, schizophrenia, etc (Orisakwe, 2014).

As a potent neurotoxin, lead may affect several cellular functions, including the release of neurotransmitters such as dopamine and serotonin. Lead is a systemic toxic, non-physiological, and non-biodegradable heavy metal found in the environment and used to produce lead-acid batteries, gasoline, water pipes, paint, solder, ceramic glazes, ammunition, etc (Omeiza *et al.*, 2021). Lead can enter different body tissues through dust inhalation, food consumption, and exposure to lead-containing materials (Dignam *et al.*, 2019). Despite extensive studies on the neurological consequences of lead and its compounds, such as lead acetate and lead chloride (PbCl<sub>2</sub>), the nontoxic exposure threshold for lead is still unknown (Omeiza *et al.*, 2021). Lead enters the systemic circulation through the intestinal mucosa and is distributed in different tissues and organs of the body. It passes through the blood-brain barrier and accumulates in the brain tissue and causes neurological disorders (Gąssowska *et al.*, 2016; Li *et*

*al.*, 2016). Lead can disrupt membrane ion channels and signaling molecules critical to neurotransmission, as well as inhibit endogenous antioxidants, thereby increasing the production of reactive oxygen/nitrogen species, and leading to oxidative stress (Bhat *et al.*, 2015; Ayuso & Foley, 2020).

Medicinal plants can be a sufficient source of new, safe, and effective treatments, especially since many of them show fewer side effects. Several medicinal plants have been studied for depression and anxiety disorders and have shown positive results (Es-Safi *et al.*, 2021).

Parsley (*Petroselinum crispum*) belongs to the Apiaceae family which is used in the food, pharmaceutical, perfume, and cosmetic industries (Lopez *et al.*, 1999). Parsley is scattered in all parts of Iran and is used in traditional medicine to treat many diseases (Ozsoy-Sacan *et al.*, 2006). Parsley has been used as a traditional medicine to treat hyperuricemia and gout in Iran. Phytochemical analysis of this plant has revealed the presence of several classes of flavonoids (Fejes *et al.*, 2000). Flavonols (kaempferol and quercetin) and flavones (apigenin and luteolin), which are found as glycosides in nature, are the main flavonoids in parsley (Peterson *et al.*, 2006). Kaempferol and quercetin, which belong to the flavonol group, have a wide range of biochemical and medicinal effects and are recommended as chemical preventive agents or food supplements. The dominant mechanism of their biological actions is due to antioxidant activity, enzyme inhibition, and free radical scavenging capacity. Therefore, it is hypothesized that the protective and therapeutic effects of parsley may be due to its flavonol compounds (Haidari *et al.*, 2011). Also, studies show that plants of the Apiaceae family have anti-depressant and anti-anxiety activities (Es-Safi *et al.*, 2021). This study aimed to investigate the protective effect of the hydroalcoholic extract of parsley leaves (HEPL) on anxiety in rats treated with lead acetate.

## Materials and Methods

### Animals

In this experimental study, 60 Wistar rats with an age range of 2.5-3 months and an approximate weight of 180±20 grams were used. The animals were obtained from the Laboratory Animal Breeding Center of the Islamic Azad University, Kazerun Branch, and were kept in polycarbonate cages with

dimensions of 30×25×15 cm with a steel mesh roof. Animals were kept under standard conditions of 12 hours light/dark, temperature 22±2°C, and humidity 70%. Rat's food was obtained from Pars Animal and Poultry Company, Iran, and full access to food and water was provided throughout the study. The ethical principles of working with laboratory animals in this study were approved by the ethics committee of the Islamic Azad University of Kazerun under the number IR.IAU.KAU.REC.1400.062.

### Animal grouping and study design

Sixty rats were randomly grouped into 6 equal groups after weighing. In the control group, the animals did not receive any drug treatment. In the sham group, the animals received 1 ml of distilled water daily as a lead acetate solvent by intraperitoneal injection. Lead group animals received lead acetate at a dose of 20 mg/kg and HEPL200 group animals received HEPL at a dose of 200 mg/kg intraperitoneally. Animals in the Lead+HEPL100 group received 20 mg/kg lead acetate intraperitoneally 2 hours before receiving HEPL at a dose of 100 mg/kg intraperitoneally. Lead+HEPL200 group animals received 20 mg/kg of lead acetate intraperitoneally 2 hours before receiving HEPL at a dose of 100 mg/kg intraperitoneally. Sixty rats were randomly grouped into 6 equal groups after weighing. In the control group, the animals did not receive any drug treatment. In the sham group, the animals received 1 ml of distilled water daily as a lead acetate solvent by intraperitoneal injection. Lead group animals received lead acetate at a dose of 20 mg/kg and HEPL200 group animals received HEPL at a dose of 200 mg/kg intraperitoneally. Animals in the Lead+HEPL100 group received 20 mg/kg lead acetate intraperitoneally 2 hours before receiving HEPL at a dose of 100 mg/kg intraperitoneally 2 hours before intraperitoneal administration of HEPL at a dose of 200 mg/kg. The length of the study period in all groups was 21 days. At the end of the study, the open box test was used to check anxiety behaviors. The parameters used in this test included Grooming (cleaning the body, face, and legs), Rearing (standing on two rear legs), and Crossing (crossing the lines).

### Preparation method of parsley leaves extract

Parsley plants were bought from farms around Kazerun City. Then, the leaves and stems of the plant were separated and dried, and powdered by scientific method. The resulting powder was placed in a

percolator and 70% ethanol was added to it. The resulting mixture was collected as HEPL in a container in the form of a drop for 24 hours. After 24 hours, diluted HEPL was prepared and then dried and concentrated using a rotary machine (Ozsoy-Sacan *et al.*, 2006). The determination of the dose of HEPL for this study was based on previous studies (Takrooni *et al.*, 2019).

### Behavioral study (Open Field Test)

This test was used to measure anxiety-like behavior (Carola *et al.*, 2002). The equipment for this test was a square board with wooden walls measuring 40 x 50 x 50 cm. Each rat was individually placed in the center of one of the open sections. Then the animal started to move in the chamber and for 6 minutes their activity was checked by a video camera. The parameters of this test included the amount of Grooming (cleaning the body, face and legs), Rearing (standing on two rear legs), and Crossing (crossing the lines), which were recorded and measured. After each test, the chamber was disinfected and cleaned with ethanol.

### Preparation method of lead acetate

Lead acetate (Sigma-Aldrich, USA) was prepared in powder form. At first, 0.5 grams of lead acetate powder was weighed with a scale with an accuracy of 0.001. Then, 100 ml of distilled water was added to the lead acetate powder as a solvent and the required volume was prepared. The determination of lead acetate dose for this study was based on previous studies (Al-Megrin *et al.*, 2019).

### Statistical Analysis


Statistical analysis of data was done using SPSS software (Version 20). One-way variance test and Tukey's post hoc test were used to compare between groups.  $P < 0.05$  was considered statistically significant and graphs were drawn with the help of GraphPad Prism 6 software.

### Results

Graph 1 shows the comparison of the mean and standard deviation of the amount of grooming in different groups. In the Lead, Lead+HEPL100, and Lead+HEPL200 groups, a significant decrease in the amount of grooming was observed compared to the control and sham groups ( $P < 0.05$ ). In the HEPL200

group, there was no significant difference in the amount of grooming compared to the control and sham groups ( $P>0.05$ ), but it showed a significant increase compared to the Lead group ( $P<0.05$ ). Also,


in the Lead+HEPL200 group, a significant increase in the amount of grooming was observed compared to the Lead group ( $P<0.05$ ).



Graph 1. A comparison of the mean and standard error of the amount of grooming in different groups is shown. \*: significant difference with control and sham groups ( $P<0.05$ ). \*\*: significant difference between Lead+HEPL100 and Lead+HEPL200 groups with Lead group ( $P<0.05$ ). Ctl (Control), Sham, Lead, HEPL (Hydroalcoholic extract of parsley leaves).

Graph 2 shows the comparison of the mean and standard deviation of rearing in different groups. In the Lead, Lead+HEPL100, and Lead+HEPL200 groups, a significant decrease in rearing was observed compared to the control and sham groups ( $P<0.05$ ). In the HEPL200 group, there was no significant


difference in rearing compared to the control and sham groups ( $P>0.05$ ), but it showed a significant increase compared to the Lead group ( $P<0.05$ ). Also, in the Lead+HEPL200 group, a significant increase in rearing was observed compared to the Lead group ( $P<0.05$ ).



Graph 2. A comparison of mean and standard error of the rearing rate in different groups is shown. \*: significant difference with control and sham groups ( $P<0.05$ ). \*\*: significant difference between Lead+HEPL100 and Lead+HEPL200 groups with Lead group ( $P<0.05$ ). Ctl (Control), Sham, Lead, HEPL (Hydroalcoholic extract of parsley leaves).

Graph 3 shows the comparison of the average and standard deviation of the amount of Crossing in different groups. In the Lead group, a significant decrease in the amount of crossing was observed compared to the control and sham groups ( $P<0.05$ ). In Lead+HEPL100, Lead+HEPL200, and HEPL200

groups, there was no significant difference in the amount of crossing compared to the control and sham groups ( $P>0.05$ ), but in the Lead+HEPL200 and HEPL200 groups, there was shown a significant increase compared to the Lead group ( $P<0.05$ )



Graph 3. A comparison of the mean and standard error of the amount of crossing in different groups is shown. \*: significant difference with control and sham groups ( $P<0.05$ ). \*\*: significant difference between Lead+HEPL100 and Lead+HEPL200 groups with Lead group ( $P<0.05$ ). Ctl (Control), Sham, Lead, HEPL (Hydroalcoholic extract of parsley leaves).

## Discussion

The results of this study showed that administration of lead for 21 days can cause anxiety behaviors in male rats. Previous studies show that exposure of rats to lead during fetal, infancy, and later can increase anxiety (Bochani *et al.*, 2011). The mechanism of action of lead may involve disruption of neurochemistry such as brain monoamine neurotransmission. Lead exposure is known to disrupt catecholaminergic systems, and depression and anxiety disorders are strongly associated with disruptions in these systems. Animal studies show that chronic exposure to lead can reduce serotonergic activity in several brain regions, including the nucleus accumbens, frontal cortex, and brainstem (Bouchard *et al.*, 2009). Lead toxicity is associated with brain oxidative stress, neuroinflammation, and excitotoxicity. Currently, the interaction between these factors is considered a characteristic of structural and functional neurological disorders in the brain after exposure to lead.

The hippocampus, an important neurobiological region for cognitive processes, is believed to be

central to the harmful effects of lead (McNamara & Skelton, 1993; Nehru & Sidhu, 2002). This study reports that exposure to lead impairs learning, induces anxiety, and induces depression-like symptoms in lead-exposed male Wistar rats (Gonenc *et al.*, 2005; Soleimani *et al.*, 2016). An imbalance between the production of reactive oxygen species (ROS) and the ability of antioxidant systems to easily detoxify these reactive intermediates leads to oxidative stress. High oxidative stress is involved in a wide range of neurological disorders (Gelderman *et al.*, 2007; Hwang, 2013). Lead damages cellular components by increasing oxidative stress because it directly disrupts enzyme activity, inactivates or depletes sulfhydryl pools of antioxidants, and increases free radical damage through direct ROS formation. (Ercal *et al.*, 2001; Patrick, 2006). Anxiety expression involves the coordinated activity of multiple brain pathways involving different neurotransmitters, all of which interact and are regulated by local and remote synaptic relays. The role of the inhibitory neurotransmitter GABA has long been considered central to the regulation of anxiety, and this



neurotransmitter system is the target of drugs used to treat anxiety disorders (Lydiard, 2003). Studies show that lead can decrease GABA levels and also affect GABA receptors through channel modification (Wirbisky et al., 2014).

Phenolic compounds are a very diverse group of plant secondary metabolites to protect cells against oxidative stress caused by free radical species (Costa et al., 2009; Oliveira et al., 2009). From a therapeutic point of view, the parsley plant shows a wide range of medicinal properties, including liver and brain protection (Farzaei et al., 2013; Sęczyk et al., 2016; Rahmat et al., 2019). Daily feeding of pregnant rats with crude extract of parsley has been shown to protect the neonatal brain and balance the deleterious changes in behavioral status, neurotransmitters, and oxidative stress caused by exposure to lead and cadmium during pregnancy (Allam et al., 2016). The effectiveness of an antioxidant can be exerted in various ways, such as scavenging free radicals, decomposing free radicals, as well as chelating metal ions (Zheng et al., 2019). Studies show that the phenolic extract of parsley has antioxidant activity (Es-safi et al., 2021). Recent studies also show the connection between mood disorders and oxidative stress (Nunes et al., 2018; Bonifácio et al., 2021) and psychological stress (Eick et al., 2018; McAllister et al., 2019). Therefore, this issue can open new ways to prevent or manage anxiety and depression about the potential use of antioxidants. Desrumaux et al, found that the lack of vitamin E as an antioxidant in the mouse brain increased the level of key signs of oxidative stress and anxiety-provoking behaviors (Desrumaux et al., 2005). Polyphenols, including flavonoids and phenolic acids, are known for their strong antioxidant effects (Mechchate et al., 2021; Fratianni et al., 2019). Phytochemical screening of parsley has revealed the presence of some compounds such as flavonoids, carotenoids, ascorbic acid, and tocopherol. These components of parsley leaves destroy superoxide anions and hydroxyl radicals in laboratory conditions. Parsley juice supplementation for 50 days has been shown to partially prevent cadmium toxicity and significantly improve physical balance, motor coordination, muscle strength, and brain neurotransmitter levels in cadmium-treated infants. (Allam et al., 2016).

Among other important compounds of the parsley plant is apigenin, which shows anti-anxiety activity

almost without side effects (Jäger & Saaby, 2011). Using these secondary phytochemical metabolites as a way to avoid and manage anxiety and depression may be a promising strategy (Amaghnoouje et al., 2020). A study conducted by Akıncı et al, showed that parsley minimizes gastric damage caused by stress when taken orally by supporting the antioxidant defense system of cells, which is reflected in increasing the average level of tissue glutathione, superoxide dismutase, and catalase activities is successful (Akıncı et al., 2017).

The open-box test is used to assess the animal's emotional state. Animals that are separated from their cages and placed in a different environment often show distress and anxiety by causing changes in all or some parameters such as a decrease in ambulatory activities, exploration and immobility, and an increase in grooming behavior. (Sturman et al., 2018). It has been shown that parsley plant extract at a dose of 100 mg/kg can produce very significant anti-anxiety effects in rats (Es-safi et al., 2021), which is consistent with the results of our study. One of the limitations of this study is the non-use of other anxiety tests such as the elevated plus maze test.

## Conclusion

The results of this study showed that administration of lead at a dose of 20 mg/kg in male rats causes anxiety behaviors. However, administration of HEPL at doses of 100 and 200 mg/kg reduced anxiety behaviors, especially at a dose of 200 mg/kg in male rats treated with lead acetate. Therefore, it is suggested to consider parsley extract as a supplement for people who are exposed to lead acetate contamination.

## Acknowledgments

The authors would like to thank all the people who helped us in conducting this study.

## Conflict of interest

There is no conflict of interest between the authors.

## References

- Akıncı A., Eşrefoğlu M., Taşlıdere E., and Ateş B. Petroselinum crispum is effective in reducing stress-induced gastric oxidative damage. *Balk Med J*, 2017; 34: 53.

- Allam AA., Maodaa SN., Abo-Eleneen R. and Ajarem J. Protective effect of parsley juice (*Petroselinum crispum*, Apiaceae) against cadmium deleterious changes in the developed albino mice newborns (*Mus musculus*) brain. *Oxid Med Cell Longev*, 2016; 2016: 2646840.
- Al-Megrin WA., Alkhuriji AF., Yousef AOS., Metwally DM., Habotta OA., Kassab RB., et al. The antagonistic efficacy of luteolin against lead acetate exposure associated with hepatotoxicity is mediated via antioxidant, anti-inflammatory, and anti-apoptotic activities. *Antioxidants (Basel)*, 2019; 9(1): 10. doi: 10.3390/antiox9010010.
- Amaghnoije A., Mechchate H., Es-Safi I., Boukhira S., Aliqahtani AS., Noman OM., et al. Subacute assessment of the toxicity and antidepressant-like effects of *origanum majorana* L. polyphenols in swiss albino mice. *Molecules*, 2020; 25: 5653.
- Ayuso RA. & Foley NK. Surface topography, mineralogy, and Pb isotope survey of wheel weights and solder: Source of metal contaminants of roadways and water systems. *J Geochem Explor*, 2020; 212: 106493. 10.1016/j.gexplo.2020.106493.
- Bhat AH., Dar KB., Anees S., Zargar MA., Masood A. and Sofi MA. Oxidative stress, mitochondrial dysfunction and neurodegenerative diseases; a mechanistic insight. *Biomed Pharmacother*, 2015; 74: 101-110. 10.1016/j.biopha.2015.07.025.
- Bochani A., Komaki A. and Nasri S. Effects of lead (Pb) exposure on anxiety in rats. *J Birjand Univ Med Sci*, 2011; 18(4): 257-264.
- Bonifácio KL., Barbosa DS., Moreira EG., Coneglian CF., Vargas HO., Nunes SOV., et al. Increased nitro-oxidative stress toxicity as a major determinant of increased blood pressure in mood disorders. *J Affect Disord*, 2021; 278: 226-238.
- Bouchard MF., Bellinger DC., Weuve J., Matthews-Bellinger J., Gilman SE., Wright RO., et al. Blood lead levels and major depressive disorder, panic disorder, and generalized anxiety disorder in US young adults. *Arch Gen Psychiatry*, 2009; 66(12): 1313-9. doi: 10.1001/archgenpsychiatry.2009.164.
- Carola V., D'Olimpio F., Brunamonti E., Mangia F. and Renzi P. Evaluation of the elevated plus-maze and openfield tests for the assessment of anxiety-related behavior in inbred mice. *Behav Brain Res*, 2002; 134: 49-57.
- Costa RM., Magalhães AS., Pereira JA., Andrade PB., Valentão P., Carvalho M., et al. Evaluation of free radical scavenging and antihemolytic activities of quince (*Cydonia oblonga*) leaf: a comparative study with green tea (*Camellia sinensis*) *Food Chem Toxicol*, 2009; 47: 860.
- Desrumaux C., Risold PY., Schroeder H., Deckert V., Masson D., Athias, A., et al. Phospholipid transfer protein (PLTP) deficiency reduces brain vitamin E content and increases anxiety in mice. *FASEB J*, 2005; 19: 1-16.
- Dignam T., Kaufmann RB., LeSturgeon L. and Brown MJ. Control of lead sources in the united states, 1970-2017: public health progress and current challenges to eliminating lead exposure. *J Public Health Manag Pract*, 2019; 25 Suppl 1, Lead Poisoning Prevention (Suppl 1 LEAD POISONING PREVENTION):S13-S22. doi: 10.1097/PHH.0000000000000889.
- Eick SM., Barrett ES., Van 't Erve TJ., Nguyen RH., Bush NR., Milne G., et al. Association between prenatal psychological stress and oxidative stress during pregnancy. *Paediatr Perinat Epidemiol*, 2018; 32: 318-326.
- Es-Safi I., Mechchate H., Amaghnoije A., Kamaly OMA., Jawhari FZ., Imtara H., et al. The potential of parsley polyphenols and their antioxidant capacity to help in the treatment of depression and anxiety: an in vivo subacute study. *Molecules*, 2021; 26(7): 2009. doi: 10.3390/molecules26072009.
- Es-safi I., Mechchate H., Amaghnoije A., Kamaly OMA., Jawhari FZ., Imtara H., et al. The potential of parsley polyphenols and their antioxidant capacity to help in the treatment of depression and anxiety: an in vivo subacute study. *Molecules*, 2021; 26(7): 2009. doi.org/10.3390/molecules26072009.

- Farzaei MH., Abbasabadi Z., Reza M., Ardekani S. and Rahimi R. Parsley: a review of ethnopharmacology, phytochemistry and biological activities. *J Traditional Chin Med*, 2013; 33: 815-826.
- Fejes SZ., Blazovics A., Lemberkovics E., Petri G., Szoke E. and Kery A. Free radical scavenging and membrane protective effects of methanol extracts from *Anthriscus cerefolium* (Hoffm) L. and *Petroselinum crispum* (Mill) Nym. Ex A W Hill *Phytother Res*, 2000; 14: 362-365.
- Fratianni F., Cozzolino A., De Feo V., Coppola R., Ombra MN. and Nazzaro F. Polyphenols, Antioxidant, antibacterial, and biofilm inhibitory activities of peel and pulp of citrus medica l., citrus bergamia, and citrus medica cv. salò cultivated in southern Italy. *Molecules*, 2019; 24: 4577.
- Gale CK. and Millichamp J. Generalised anxiety disorder. *BMJ Clin Evid*, 2011; 2011:1002.
- Gąssowska M., Baranowska-Bosiacka I., Moczyłowska J., Frontczak-Baniewicz M., Gewartowska M., Strużyńska L., et al. Perinatal exposure to lead (Pb) induces ultrastructural and molecular alterations in synapses of rat offspring. *Toxicology*, 2016; 373:13-29. doi: 10.1016/j.tox.2016.10.014.
- Ghasemzadeh Dehkordi S., Doulah A. and Rafieirad M. The effect of four weeks of exercise and oleuropein supplementation on anxiety and depression behaviors in an animal model of Parkinson's disease. *Applied Biology*, 2022; 35(1): 180-195. doi: 10.22051/jab.2021.36815.1432.
- Gonenc S., Uysal N., Acikgoz O., Kayatekin BM., Sonmez A., Kiray M., et al. Effects of melatonin on oxidative stress and spatial memory impairment induced by acute ethanol treatment in rats. *Physiol Res*, 2005; 54(3): 341-348.
- Haidari F., Keshavarz SA., Mohammad Shahi M., Mahboob SA. and Rashidi MR. Effects of parsley (*Petroselinum crispum*) and its flavonol constituents, kaempferol and quercetin, on serum uric acid levels, biomarkers of oxidative stress and liver xanthine oxidoreductase activity in oxonate-induced hyperuricemic rats. *Iran J Pharm Res*, 2011; 10(4): 811-9.
- Jäger AK. and Saaby L. Flavonoids and the CNS. *Molecules*, 2011; 16: 1471-1485.
- Li X., Ye F., Li L., Chang W., Wu X. and Chen J. The role of HO-1 in protection against lead-induced neurotoxicity. *Neurotoxicology*, 2016; 52:1-11. doi: 10.1016/j.neuro.2015.10.015.
- Lopez MG., Sanchez-Mendoza IR. and Ochoa-Alejo N. Comparative study of volatile components and fatty acids of plants and in-vitro cultures of parsley *Petroselinum crispum* (Mill) nym ex hill. *J Agric Food Chem*, 1999; 47: 3292-3296.
- Lydiard RB. The role of GABA in anxiety disorders. *J Clin Psychiatry*, 2003; 64(Suppl 3): 21-27.
- McAllister MJ., Basham SA., Waldman HS., Smith JW., Mettler JA., Butawan M.B., et al. Effects of Psychological Stress during Exercise on Markers of Oxidative Stress in Young Healthy, Trained Men. *Physiol Behav*, 2019; 198: 90-95.
- McNamara RK. and Skelton RW. The neuropharmacological and neurochemical basis of place learning in the Morris water maze. *Brain Res Rev*, 1993; 18(1): 33-49. doi: 10.1016/0165-0173(93)90006-L.
- Mechchate H., Es-safi I., Amaghnouje A., Boukhira S., Alotaibi AA., Al-zharani M., et al. Antioxidant, anti-inflammatory and antidiabetic properties of LC-MS/MS identified polyphenols from coriander seeds. *Molecules*, 2021; 26: 487.
- Nehru B. and Sidhu P. Neurotoxic effects of differential doses of lead on rat brain followed by recovery. *JTEMIN*, 2002; 15(3): 131-140. doi: 10.1002/jtra.10009.
- Nunes CS., Maes M., Roomruangwong C., Moraes JB., Bonifacio KL., Vargas HO., et al. Lowered quality of life in mood disorders is associated with increased neuro-oxidative stress and basal thyroid-stimulating hormone levels and use of anticonvulsant mood stabilizers. *J Eval Clin Pract*, 2018; 24: 869-878.



- Oliveira AP., Valenao P., Pereira JA., Silva BM., Tavares F. and Andrade PB. Ficus carica L.: metabolic and biological screening. Food Chem Toxicol, 2009; 47: 2841-2846.
- Omeiza NA., Abdulrahim HA., Alagbonsi AI., Ezurike PU., Soluoku TK., Isiabor H., et al. Melatonin salvages lead-induced neuro-cognitive shutdown, anxiety, and depressive-like symptoms via oxido-inflammatory and cholinergic mechanisms. Brain Behav, 2021; 11(8): e2227. doi: 10.1002/brb3.2227.
- Orisakwe OE. The role of lead and cadmium in psychiatry. N Am J Med Sci, 2014; 6(8): 370-6. doi: 10.4103/1947-2714.139283.
- Ozsoy-Sacan O., Yanardag R., Orak H., Ozgey Y., Yarat A. and Tunali T. Effects of parsley (*Petroselinum crispum*) extract versus glibornuride on the liver of streptozotocin-induced diabetic rats. J Ethnopharmacol, 2006; 104: 175-181.
- Ozsoy-Sacan O., Yanardag R., Orak H., Ozgey Y., Yarat A. and Tunali T. Effects of parsley (*Petroselinum crispum*) extract versus glibornuride on the liver of streptozotocin-induced diabetic rats. J Ethnopharmacol, 2006; 104(1-2): 175-81.
- Peterson S., Lampe JW., Bammler TK., Gross-Steinmeyer K. and Eaton DL. Apiaceous vegetable constituents inhibit human cytochrome P-450 1A2 (hCYP1A2) activity and hCYP1A2-mediated mutagenicity of aflatoxin B1. Food Chem Toxicol, 2006; 44: 1474-1484.
- Rahmat A., Ahmad NSS. and Ramli NS. Parsley (*Petroselinum crispum*) supplementation attenuates serum uric acid level and improves liver and kidney structures in oxonate-induced hyperuricemic rats. Orient Pharm Exp Med, 2019; 19: 393-401.
- Sęczyk Ł., Świeca M., Gawlik-Dziki U., Luty M. and Czyż J. Effect of fortification with parsley (*Petroselinum crispum* Mill.) leaves on the nutraceutical and nutritional quality of wheat pasta. Food Chem, 2016; 190: 419-428.
- Soleimani E., Goudarzi I., Abrari K. and Lashkarbolouki T. The combined effects of developmental lead and ethanol exposure on hippocampus dependent spatial learning and memory in rats: Role of oxidative stress. FCT, 2016; 96: 263-272. doi: 10.1016/j.fct.2016.07.009
- Sturman O., Germain PL. and Bohacek J. Exploratory rearing: a context-and stress-sensitive behavior recorded in the open-field test. Stress, 2018; 21: 443-452.
- Takrooni WA., Sharaf IA. and Abdul Majid NA. Assessment of the potential role of parsley (*Petroselinum Crispum*) leaves extract in ameliorating cyclosporin a- induced nephrotoxicity in rats. Int J Pharm Res Allied Sci, 2019; 8(2): 118-128.
- Weiss B., Cory-Slechta D., Gilbert SG., Mergler D., Miller E., Miller C., et al. The new tapestry of risk assessment. Neurotoxicology, 2008; 29: 883-90.
- Wirbisky SE., Weber GJ., Lee JW., Cannon JR. and Freeman JL. Novel dose-dependent alterations in excitatory GABA during embryonic development associated with lead (Pb) neurotoxicity. Toxicol Lett, 2014; 229(1): 1-8. doi: 10.1016/j.toxlet.2014.05.016.
- Zheng X., Cheng Y., Chen Y., Yue Y., Li Y., Xia S., et al. Ferulic acid improves depressive-like behavior in prenatally-stressed offspring rats via anti-inflammatory activity and hpa axis. Int J Mol Sci, 2019; 20: 493.



## بررسی تاثیر عصاره هیدروالکلی برگ گیاه جعفری (*Petroselinum crispum*) بر اضطراب در موشهای صحرایی تیمار شده با استات سرب

فاطمه بسطام پور<sup>۱</sup>، سید ابراهیم حسینی<sup>۲،\*</sup>، مهرداد شریعتی<sup>۱</sup>، مختار مختاری<sup>۱</sup>

<sup>۱</sup>گروه زیست شناسی، واحد کازرون، دانشگاه آزاد اسلامی، کازرون، ایران

<sup>۲</sup>گروه زیست شناسی، دانشکده علوم، موسسه آموزش عالی زند، شیراز، ایران

تاریخ دریافت: ۱۴۰۱/۱۰/۲۷ اصلاح نهایی: ۱۴۰۱/۱۲/۰۱ تاریخ پذیرش: ۱۴۰۱/۱۲/۱۰

### چکیده

**زمینه و هدف:** فلزات سنگین مانند سرب (Lead) می‌توانند تأثیرات منفی بر سیستم عصبی داشته باشند. اضطراب یکی از شایعترین اختلالات عصبی است که می‌تواند تحت تأثیر سرب القا شود. هدف از انجام این مطالعه بررسی تأثیر حفاظتی عصاره هیدروالکلی برگ گیاه جعفری (*Petroselinum crispum*; HEPL) بر اضطراب در موشهای صحرایی تیمار شده با استات سرب بود.

**مواد و روش‌ها:** موشهای صحرایی نر به ۶ گروه ۱۰ تایی تقسیم شدند. گروه کنترل تیمار دارویی دریافت نکرد اما حیوانات گروه شم ۱ میلی‌لیتر آب مقطر به عنوان حلال استات سرب به صورت تزریق درون صفاقی دریافت کردند. حیوانات گروه Lead، استات سرب را با دوز ۲۰ mg/kg و حیوانات گروه HEPL، HEPL200 با دوز ۲۰۰ mg/kg به صورت تزریق درون صفاقی دریافت کردند. حیوانات گروه‌های Lead+HEPL100 و Lead+HEPL200 به ترتیب ۲ ساعت قبل از دریافت درون صفاقی HEPL با دوز ۱۰۰ و ۲۰۰ mg/kg استات سرب با دوز ۲۰ mg/kg به صورت تزریق درون صفاقی دریافت کردند. دوره مطالعه در تمام گروه‌ها ۲۱ روز در نظر گرفته شد. در انتهای مطالعه از آزمون جعبه باز برای اندازه‌گیری رفتار اضطرابی شامل پارامترهای Grooming (تمیز کردن بدن، صورت و پاها)، Rearing (ایستادن روی دو پای عقبی) و Crossing (عبور از خطوط) استفاده شد. تجزیه و تحلیل آماری با استفاده از آنالیز واریانس یک طرفه و تست تعقیب Tukey انجام شد.

**یافته‌ها:** نتایج این مطالعه نشان داد که در گروه Lead کاهش معنادار تعداد Rearing، Grooming و Crossing در مقایسه با گروه‌های کنترل و شاهد مشاهده گردید ( $P < 0.05$ ). در گروه Lead+HEPL200 افزایش معنادار تعداد Rearing، Grooming و Crossing در مقایسه با گروه Lead مشاهده گردید ( $P < 0.05$ ).

**نتیجه‌گیری:** سرب می‌تواند موجب ایجاد رفتارهای اضطرابی در موشهای صحرایی نر شود. با این حال تجویز HEPL می‌تواند موجب بهبود رفتارهای اضطرابی در موشهای صحرایی نر تیمار شده با استات سرب شود.

**واژه‌های کلیدی:** سرب، عصاره هیدروالکلی، جعفری، اضطراب، موش صحرایی

فاطمه بسطام پور، سید ابراهیم حسینی، مهرداد شریعتی، مختار مختاری. بررسی تأثیر عصاره هیدروالکلی برگ گیاه جعفری (*Petroselinum crispum*) بر اضطراب در موشهای صحرایی تیمار شده با استات سرب. مجله طب دامپزشکی جایگزین. ۱۴۰۲؛ ۶(۱۶): ۹۲۹-۹۳۸.

\* نویسنده مسئول: گروه زیست شناسی، دانشکده علوم، موسسه آموزش عالی زند، شیراز، ایران.