





## Effect of Human Chorionic Gonadotropin and Flunixin Meglumine on Pregnancy Rate in Dairy Cows

Navid Aarabi\*, Mohammad Mehdi Hadipour

Department of Clinical Sciences, Faculty of Veterinary Medicine, Kazerun Branch, Islamic Azad University, Kazerun, Iran

Received: 09/Nov/2023 Revised: 21/Dec/2023 Accepted: 29/Dec/2023

#### Abstract

**Background and aim:** Growing infertility and a reduction in pregnancy rates are two of the most challenging issues facing modern dairy cows. Our objectives were to determine pregnancy rates, plasma progesterone concentrations, and corpus luteum (CL) numbers in high-producing dairy cows during two distinct periods (cool and warm seasons). This evaluation followed the administration of: 1) human chorionic gonadotropin (hCG) on day 5 after AI, 2) a combined treatment with hCG on day 5 and flunixin meglumine between days 14 and 16 after AI, and 3) flunixin meglumine alone between days 14 and 16 after AI.

**Materials and Methods:** Cows were subjected to pre-synch programming in two experiments (cool period, n = 396) and warm period, n = 181), respectively. Environmental data in the warm period indicated that cows experienced high heat stress during the trial in the warm period (THI = 85-92). Cows in both experiments were randomly divided into four groups: Group 1 received hCG, and Group 2 was given hCG on day 5 and FM on days 14-16, Group 3 was treated with FM on days 14-16, and the untreated group was the control group. Ovarian structures and serum progesterone levels were determined on days 5 and 12 after AI, respectively. Ultrasonography was used to evaluate the pregnant status of cows on days 28-30.

**Results:** There was a significant difference in p4 concentrations 12 days after IA between groups in the cool period, in both the hCG and hCG+FM groups (p = 0.001). But the discrepancy in p4 was not observed between groups in the warm period (p = 0.7). The increase in CL for cows treated with hCG compared with control cows was greater during the cool (1.7 vs. 1.1) period compared to the warm (1.1 vs. 1.3) period (p = 0.001). In the warm season, no significant difference in pregnancy rate on day 30 between the four groups was obtained.


Conclusion: Our results indicated that administration of flunixin meglumine did not improve pregnancy rates in dairy cows during either the warm or cool seasons. Conversely, treatment with hCG on day 5 after artificial insemination induced accessory corpora lutea, increased plasma progesterone concentration, and improved pregnancy rates in dairy cows during the cool period only. None of these effects were observed during the warm period.

**Keywords:** hCG, Flunixin meglumine, Progesterone, Pregnancy, Heat stress

Cite this article as: Navid Aarabi, Mohammad Mehdi Hadipour. Effect of human chorionic gonadotropin and flunixin meglumine on pregnancy rate in dairy cows. J Altern Vet Med. 2024; 7(20): 1183-1194.

## \* Corresponding Author

Department of Clinical Sciences, Faculty of Veterinary Medicine, Kazerun Branch, Islamic Azad University, Kazerun, Iran.





### Introduction

Embryonic death reduces bovine fertility and causes significant economic loss in the dairy industry (Kidder et al., 1954; Diskin & Sree Nan, 1980), which has been attributed to a variety of factors such as herd size, management, metabolic, and disease states, as well as changes in physiology associated with increased milk production (Lopez et al., 2004; Wiltbank et al., 2006, Caraviello, et al., 2006). One of the most critical variables limiting dairy cow reproductive success is estrus detection effectiveness. Controlling follicular dynamics and luteal life span is one strategy for managing the reduction in reproductive performance of dairy cattle (Lucy, 2001). Synchronized cows usually have a small ovulatory follicle at the time of insemination; furthermore, it would result in the production of smaller corpus luteam that may produce lower progesterone concentrations that are optimal for fertility (Vasconcelos et al., 2001; Perry et al., 2005; Gaja et al., 2008). This problem deteriorates due to the high metabolism of P4 that is characteristic of high-producing dairy cows (Sangsritavong et al., 2002; Wiltbank et al., 2006). Progesterone plays a vital role in maintaining pregnancy and triggers the production of an extensive range of endometrial compounds, growth factors, immunosuppressive agents, enzymes, ions, and steroids (Geisert et al., 1992; Forde et al., 2011). One study suggests that combined injections of human chorionic gonadotropin (hCG) and flunixin meglumine could potentially improve pregnancy rates in cows experiencing repeat breeding (Jawad et al., 2021).

A study showed that the use of progesterone supplementation from days 5–9 following insemination can increase both embryonic development and the secretion of interferon τ (Mann et al., 2006). Findings by Satterfield showed that administration of progesterone 36 hours after mating results in increasing the blastocyst diameter on day 9 in sheep (Satterfield et al., 2006). Several studies have shown that low progesterone concentrations after post-insemination reduce fertility in dairy cattle (Stevenson et al., 1993, Gumen & Wiltbank, 2005; Diskin et al., 2006). To tackle this problem, numerous increase peripheral therapies are available to progesterone concentration following AI. encompassing inducing accessory CL or supplementing progesterone directly (Inskeep, 2004;

Stevenson et al., 2007). hCG is one of the most often used methods to induce CL; administration of hCG during the early phase results in new CL, which leads to an increase in progesterone concentration (Stevenson et al., 2007; Santos et al., 2001). Maternal recognition of pregnancy initiates approximately 14 days after breeding (Bazer et al., 1991; Roberts et al., 1996; Thatcher et al., 1994). If interferon  $\tau$  is not secreted sufficiently, the uterus produces PGF2α, which causes the corpus lutea to regress, decrease progesterone secretion, and reduce the pregnancy rate (Roberts et al., 1996; Roberts et al., 1992; Thatcher et al., 2001). Flunixin meglumine (FM) is an antiinflammatory agent that inhibits cyclooxygenase, preventing the conversion of arachidonic acid to PGF2α (Anderson et al., 1990; Odens et al., 1995).

One study found that giving FM on day 14 after AI results in a 10% increase in pregnancy rates in beef cattle due to a decrease in serum PGFM (Merrill et al., 2005). FM treatment has been shown to inhibit PGf2 secretion in beef and dairy cattle for at least 24 hours (Guilbault et al., 1987). Heat stress reduces pregnancy rates due to its detrimental effects on follicle development, corpus lutea function, and hormone profiles, leading to early embryo death (Hansen & Arechiga, 1999; Badinga et al., 1985; Barker et al., 1994).

According to one research, fertilization rates are normal, but mortality occurs between conception and maternal pregnancy identification (Thatcher & Collier, 1986; Wise et al., 1988). It has been observed that heat stress has an effect on the oocyte by destroying it, affecting protein synthesis throughout the morula and blastocyst stages, and perhaps decreasing endometrial function (Hansen et al., 2007). Many different methods including as the administration of HCG, GnRH, CIDR, and NSAIDs, have been used to minimize the effect of heat stress on reproductive performance. The use of hCG to increase pregnancy rates under heat stress is still controversial. Some studies found a favorable impact on pregnancy rates, while others did not (Dunlap & Vincent, 1971). The current study intended to determine if administering hCG and FM on days 5 and 13, respectively, following AI would enhance conception rates in warm and cool seasons. In addition, the effect of hCG on CL number and plasma progesterone concentration was evaluated.

## **Materials and Methods**

The study was conducted on 2500 lactating Holstein-Friesian cows from a single, well-managed dairy herd. The cows were kept in open stalls and fed a total mixed ration (TMR) three times a day. The TMR was formulated to meet or exceed requirements for lactating cows milked three times in a day, according to the NRC. Voluntary waiting Period was around 50-60 days postpartum, the parity of the cows ranged from 2 to 4, and the body condition score was determined on the basis of scales 1-5 (NRC, 2001). Cows undergoing an abnormal puerperium, such as dystocia, twinning, retained placenta, or metritis, were excluded. Cows with clinical conditions detected during the course of the study, such as mastitis, lameness, digestive disorders, abnormal genital discharges, and pathological abnormalities of the reproductive tract detectable on palpation per rectum, were also withdrawn from the program. All the animals in cool (n = 396) and warm periods (n =181) were pre-synchronized with two cloprostenol (lutalyz, Pfizer, USA) IMs. Treatments were given 14 days apart, starting on day 30-35 postpartum. Twelve days following the second PGF2a treatment, the Ovsynch protocol, consisting of an injection of GnRH Co.), Aburaihan followed (Vetaroline; administration of PGF2a 7 days later, and a final injection of GnRH given 24 hours after the third PGF2a treatment, began. Cows were subjected to timed AI 16-22 hours after the last injection of GnRH. All cows were inseminated by the same technician, and semen from seven proven sires was used. The cows were randomly divided into four groups. Group-1 (cool period, n=100), (warm period, n=50) received 1500 IU hCG (IM) on day 5 after the initial AI; group-2 was treated with 100 µg FM (Flunixin meglumine, flunex, Razak) for three days (IM) between days 14 and 16 post-insemination (cool period, n=98) and (warm period, n=50); group 3 received 1500 IU hCG on day 5 and 100 µg FM for 3 days on days 14–16 after AI (cool period, n=101); (warm period, n=41); and group 4 was selected as the control with no treatments (cool period, n=97) and (warm period n=40).

Blood samples were collected via venipuncture of the median caudal vein or artery into evacuated tubes on days 5, 12 after TAI for later analysis of serum P4. The first sample was collected immediately before

administration of treatment. Upon collection, samples were immediately placed on ice, and serum was separated by centrifugation (for 10 min at  $3000 \times g$ ) and stored at -22 °C until assayed. Serum P4 concentrations on days 5, 12 after AI were determined by using a validated commercial radio immunoassay kit (Immunotech kit, Marseille, France). The intra- and inter-assay coefficients of variation (CVs) of progesterone were 6.5 and 9.0%, respectively. The sensitivity of the test was 0.05 ng/mL, and the recovery rate of the assay ranged from 85 to 110%. Ovarian structures on d 5 and 12 were scanned and mapped using transrectal ultrasonography (7.5 MHZ transrectal probe, SIUI, CT 900V, China). Ultrasound measurements of CL dimensions (length (L) and width (W)) were used to calculate luteal tissue volume. First, the radius (R) was calculated by the formula R = (L/2 + W/2)/2, and then the volume (V) by using the formula V =  $4/3*\pi*R^3$  (Vasconcelos et al., 2001). The pregnancy status of cows was estimated for four groups at days 28-30 and confirmed on days 40-45 ultrasonography.

Temperature- humanity index (THI) was calculated as per following equation:

T-H index = 0.81 T + 0.01 RH (0.99 T-14.3) + 46.3Where: T: mean air temperature in °C.

RH: mean relative humidity%.

## Statistical analysis

Data were imported into SAS version 9.2 (SAS Institute Inc., Cary, NC, USA, 2010) for statistical analysis. Descriptive statistics were done using PROC MEANS in SAS. All dependent data were checked to fit a normal distribution using the Shapiro-Wilk test (Shapiro & Wilk, 1965). Qualitative data from the 8 groups was compared by chi-square analysis (Proc FREO, SAS). Differences in corpus luteal volume and progesterone concentrations between subjects treated with hCG, hCG+FM, FM, and untreated were tested by the ANOVA procedure. To confirm the present hypothesis that hCG administration on day 5 after AI increases the number of CLs and progesterone concentration, two sample t-tests were used to compare the numbers of CL and p4 values on day 5. Compared with day 12 after AI in the control group as well as in the treatment group. Pearson correlation was used to assess the relationship

between CL volume and plasma p4 concentration on day 12 after AI in groups separately. Regression analyses were conducted according to the method of Hosmer and Lemeshow (Hosmer & Lemeshow, 1989). The selected potential risk factors were then subjected to stepwise logistic regression approaches with hierarchical backward elimination and swapping. The P-values for data inclusion and exclusion were set at 0.05 and 0.10, respectively. All variables that had been selected or retained in those stepping approaches entered the final likelihood ratio (LR), from which the final odds ratio estimates with 95% confidence intervals were derived. P-values <0.05 were considered statistically significant.

## Results

The mean ( $\pm$ SD) milk yields of the cows during the study were 39.4 $\pm$ 3.4 kg/day in the cool season and 35.4 $\pm$ 3.2 kg/day in the warm season.

In the cool season, there was a significant difference between the pregnancy rate on the day 28-30 in the treated (hCG, p = 0.03) (56%), (hCG+FM, p = 0.05) (55.1%), control (41.2%), and treated FM (40.6%) groups. Whereas, there was no significant difference between the treated (FM) and control groups. In the warm season, no significant differences in pregnancy rate on day 30 between the four groups were obtained (Figure 1 and 2).




Figure 1. Pregnancy rate between four groups on d 30 of gestation in cool period.

The formation of the accessory luteal structure was greater (hCG, p = 0.001) and (hCG+FM, p = 0.001) for cows in the treatment groups (68%, 70.4%), respectively, than for the ones in the control group (26.5%). But in the warm season, no significant difference in the number of CLs between groups was observed. The increase in CL for cows treated with hCG compared with control cows was greater during the cool (1.7 vs. 1.1) period than during the warm (1.1 vs. 1.3) period (p = 0.001). There was a positive correlation between BSC and pregnancy, as well as

the number of CL in hCG and hCG+FM - treated cows, so that the rate of pregnancy and CL number increased in cows with a BSC greater than 2.75 compared with cows with a BSC lower than 2.75. (Table 1).

The discrepancy was not obtained in p4 concentration on day 5 after IA between groups in the cool (p = 0.8) and warm periods (p = 0.6). Whereas there was a significant difference in p4 concentrations on the 12th day after IA between groups in a cool period. In both the hCG and hCG+FM groups, the amount of p4 was greater than the other groups (p = 0.001). But the




Figure 2. Pregnancy rate between groups on d 30 of gestation in warm period.

discrepancy in p4 was not observed between groups in the warm period (p = 0.7) (Table 2).

On Days 5 and 12, there were no differences in CL volumes between the third and control groups (p = 0.7). The discrepancy was observed in the CLs volume in the treatment hCG and hCG +FM groups from Days 5 to 12. The average volumes of CL were greater on Day 12 in the treated groups than the other groups (p = 0.001) (Table 2).

The significant association between CLs volume and progesterone concentrations on days 5 and 12 in both hCG and hCG +FM treated was obtained in the cool period (r = 0.4; p = 0.0001). But there was no significant correlation between serum progesterone and CL volume on days 5 and 12 in both the control and FM-treated groups (p = 0.09).

	Pregnancy status				The number of CL				
	open		Pregnant	Pregnant		1		1>	
BCS	hCG	hCG+FM	hCG	hCG+FM	hCG	hCG+FM	hCG	hCG+FM	
2.75<	87%	79.50%	13% a	20.50% a	52.20%	59.10%	26% a	22.70% a	
2.75 ≥	31.20%	34.40%	68.80% b	65.60% ь	47.80%	40.90%	74% b	77.30% в	

**Table 1.** Effect of body condition score at AI on pregnancy rate and the number of CL. a·b Data with different subscription in column significantly differ p≤ 0.05.

On Day 5, plasma progesterone concentrations were similar in the cold period for cows in the control, hCG, hCG+FM, and FM groups (P = 0.8). On Day 12, plasma progesterone concentrations in cows in the treatment groups (hCG, hCG+FM) were greater (P = 0.001) ( $13\pm3.7$  ng/mL) ( $12.7\pm3.9$ ), respectively, than in cows in the control and FM groups ( $9.7\pm2.9$ ), ( $10\pm2$ ).

The rising number of CLs and treatment during the cool period improved the odds of the pregnancy rate by 2.5 and 1.8 times, respectively. The rise in concentration of progesterone following hCG treatment increased the odds of a pregnancy by 1.1 times (Table 3).

Cool period						Warm period			
		Trea	tment		Treatment				
	hCG	hCG+FM <sup>2</sup>	FM	Control	hCG	hCG+FM	FM	Control	
Item									
Progesterone: ng/mL									
on day 5	6.1	5.7	4.9	5.2	4.8	5.5	5.8	6.1	
on day 12	13 a	12.7 a	10 ь	9,7 ь	11.3	10.4	9.8	9.6	
CL1 number	1.68 a	1.70 a	1.19 в	1.21 <sup>b</sup>	1.21	1.16	0.90	1.11	
CL volume mm <sup>3</sup>	14253±8850a	14331±12390a	9672±8680 <sup>b</sup>	9210±1781 <sup>b</sup>	9113±8343	10012±12426	9272±2212	9865±8645	

**Table2.** Effect of treatment on plasma progesterone concentration at days 5 and 12 and corpora lutea number at days. 12 after AI. <sup>a.b</sup> means with different superscript within rows are different; p< 0.05, 1-corpus lutea, 2-flunxin meglumine.

Variable	Class	Odds ratio	95% Confidence interval	P- value
Number of cl	CL > 1	2.5	1.33 - 4.92	0.01
	CL = 1	Reference		
Season	cool period	1.8	1.11 - 3.2	0.005
	warm period	Reference		
Progesterone	continuous	1.1	1.045 - 1.2	0.002

**Table 3.** Logistic regression predicting likelihood of pregnant cows Backward likelihood ratio test = 77.8, 4 df, p= 0.0001. Hosmer and lemeshow goodness – of – fit, test=3.17, 8 df, p= 0.9 the model fits.

#### Discussion

Progesterone is essential for the formation and maintenance of pregnancy. Between days 8 and 16 of pregnancy, the mortality rate is estimated to be over 40% (Diskin & Morris, 2008; Sreenan & Diskin, 1986). Several studies have found that administering hCG on day 5 after AI increases the concentration of progesterone in dairy cows, heifers, and beef cows (Stevenson et al., 2007; Santos et al., 2001; Chagas et al., 2005; Nishigai et al., 2005). The administration of hCG on the fifth day following estrus results in an increase in progesterone due to the development of accessory CLs (Schmitt et al., 1996b). Fricke and a colleague observed that a 1500 IU hCG injection on day 6 following estrus triggered ovulation and the formation of a new CL (Fricke et al., 1993). A study found that administering flunixin meglumine during embryo transfer increases pregnancy rates in both Angus cows and Holstein heifer (Tahir et al., 2021). Another study suggests that combined injections of human chorionic gonadotropin (hCG) and flunixin meglumine could potentially improve pregnancy rates in cows experiencing repeat breeding (Jawad et al., 2021).

Our data showed that administration of hCG on day 5 after AI increased the number of CL and progesterone concentrations on day 12 after AI in the

cool season, but administration of hCG on day 5 after AI did not increase the number of corpora lutea and plasma progesterone concentrations on day 12 after AI in the warm season, which is consistent with Santos's (Santos et al., 2001) study, which demonstrated that injection of hCG on day 5 after AI did not increase the number of CL and progesterone concentration in the warm period (Santos et al., 2001). Schmitt and colleagues found that inducing an accessory CL with hCG on days 5 or 6 after insemination did not increase pregnancy rates in fertile heifers and lactating dairy cows during summer heat stress (Schmitt et al., 1996b). Another study found that giving hCG to cows on day 5 following AI during the warm season had no effect on pregnancy rates (Hansen & Arechiga, 1999).

Two studies suggest that injecting flunixin meglumine into recipient beef cows undergoing embryo transfer could potentially increase their pregnancy rates (Kasimanickam *et al.*, 2018; Kasimanickam *et al.*, 2019). Administration of human chorionic gonadotropin (hCG) on days 4–7 after insemination has been shown to increase progesterone levels and improve pregnancy rates in dairy cows, as evidenced by studies demonstrating both effects (Santos *et al.*, 2001; walker *et al.*, 2005; Rensis *et al.*, 2008). Additionally, a study published reported that

administering hCG on day 5 after insemination elevated progesterone concentrations and improved pregnancy rates in both the hCG and ketoprofen groups (Singh, 2020).

Cattle embryos are susceptible to high temperatures during the first 2 to 3 days after fertilization, and they will die if exposed to high temperatures (Hansen et al., 1999). Therefore, it is unlikely that inducing an accessory CL by treatment with hCG on day 5 after AI would improve the conception rate of heat - stressed dairy cows. Wolfenson and coworkers (1995) observed that cows had a 2 to 3 days earlier emergence of the secondwave dominant follicle and decreased the number of medium-sized follicles in heat stress (Wolfenson et al., 1995). Perhaps, by d 5 after AI, the first-wave dominant follicle of some cows might have lost dominance, and that would decrease the response to hCG during the warm period. We determined a significant correlation between plasma progesterone concentration and CL volume, which agreed with earlier research that found a strong positive relationship between CL volume and plasma progesterone (Santos et al., 2001; Stevenson et al., 2007; Raja Mahendran & Sianangama, 1992). The effect of hCG on early embryo death is debatable. Some studies found that giving hCG on day 5 following AI improved the pregnancy rate (Breuel et al., 1990; Sianangama & Mahendran 1992; Singh, 2020), whereas others found no change in the pregnancy rate in lactating cows (Eduvie & Seguin, 1982, Hansel et al., 1976). Our findings show that treated cows with hCG on day 5 after AI increased pregnancy rate on day 30 after AI compared to the control group during the cool season but did not improve pregnancy rate in the warm season, which is consistent with the findings of Santos, who found that administration of hCG on day 5 after AI increased pregnancy rate on day 28 after AI during the cool season but had similar conception rates during the warm season (Santos et al., 2001). Some studies found that administering FM after AI enhanced the likelihood of pregnancy (Merrill et al., 2005) or at the time of embryo transfer (Purcell et al., 2005; Scenna et al., 2005). Treatment with FM on day 14 after AI may reduce PGf2 secretion and allow for more time for maternal recognition.

A similar result as ours was obtained in one study, which indicated that administration of GnRH on day 5 and another study showed that injections of hCG on day 6 after AI and ketoprofen on day 11 following AI improved pregnancy rates (Kraevskiy *et al.*, 2020; Rossetti *et al.*, 2011). However, it has been reported that administration of hCG on the 7<sup>th</sup> day and flunixin meglumine on the 16<sup>th</sup> day after insemination could not increase the pregnancy rate compared with the untreated group in Nelor cows (Eyyüp & Cevdet, 2023).

Research conducted found that giving FM to cows on days 13-15 following AI did not boost the conception rate when compared to the control group in both cool and warm weather. FM administration on day 13 following AI failed to improve pregnancy establishment in beef heifers and cows, according to Greay and colleagues (Geary et al., 2009). One study showed that FM-treated cows and heifers had higher pregnancy rates than untreated cows, which contradicted our findings (Merrill et al., 2005). Another study found that giving FM to cows on day 13 following AI did not boost conception rates in the warm season (Kaveh et al., 2007). Cows with low BCS are more likely to be in anestrus and have poorer fertility, according to Ferguson (Ferguson, 1991). Our findings confirmed the Santos study, which found that cows with a BCS greater than 2.75 at the time of artificial insemination had a higher pregnancy rate and number of CL than those with a BCS equal to or lower than 2.75 (Santos et al., 2001).

## Conclusion

Our data show that hCG therapy on day 5 following AI in dairy cows significantly increased serum progesterone concentrations, the number of CLs, and the pregnancy rate compared to the control group in the cold period. The administration of flunixin, meglumine in both cool and warm periods, and hCG to heat-stressed cows had no effect on plasma progesterone levels, the number of CLs, or the pregnancy rate.

## **Conflict of Interest**

None declared.

## References

Anderson KL., Neff-Davis CA., Davis LE. and Bass VD. Pharmacokinetics of flunixin meglumine in lactating cattle after single and multiple

- intramuscular and intravenous administrations. Am J Vet Res, 1990; 51: 1464-1467.
- Badinga L., Collier RI. and Wilcox CJ. Effects Thatcher of climatic and management factors on conception rate of dairy cattle in subtropical environment. J Dairy Sci, 1985; 68: 78-85.
- Barker R., Riso C., Donovan GA. and Comp Cont. low palpation pregnancy rate resulting from low conception rate in a dairy herd with adequate estrus detection intensity. Edu Part Vet, 1994; 16: 801.
- Bazer FW., Thatcher WW., Hansen PJ., Mirando MA., Ott TL. and Plante C. Physiological mechanisms of pregnancy recognition in ruminants. J Reprod Fert Suppl, 1991; 43: 39-47.
- Breuel KF., Spitzer JC., Thompson CE. and Breuel JF. First-service pregnancy rate in beef heifers as induced by human chorionic gonadotropin administration before and/or after breeding. Theriogenology, 1990; 34-36.
- Caraviello DZ., Weigel KA., Craven M., Gianola D., Cook NB., Nordlund KV., et al. Analysis of reproductive performance of lactating cows on large dairy farms using machine learning algorithms. J Dairy Sci, 2006; 89: 4703-4722.
- Chagas e Silva J. and Lopes da Costa L. Luteotropic influence of early bovine embryos and the relationship between plasma progesterone concentrations and embryo survival. Theriogenology, 2005; 64: 49-60.
- Diskin MG. and Sree Nan JM. Fertilization and embryonic mortality rates in beef heifers after artificial insemination. J Reprod Fertil, 1980; 463-468.
- Diskin MG. and Morris DG. Embryonic and early fetal losses in cattle and other ruminants. Reprod Domest Anim, 2008; 43(2): 260-267.
- Diskin MG., Murphy JJ. and Sree nan JM. Embryo survival in dairy cows managed under pastoral conditions. Anim Reprod Sci, 2006; 96: 297-311.

- Dunlap SE. and Vincent CK. Effect of sex on growth performance in pigs, 1971; J Anim Sci 32:1216.
- Eduvie LO. and Seguin BE. Corpus luteum function and pregnancy rate in lactating dairy cows given human chorionic gonadotropin at mid diestrus. Theriogenology, 1982; 17(4):
- Eyyüp Hakan U. and Cevdet P. The Effect of Human Chorionic Gonadotropin and Ketoprofen Applications on Pregnancy Rates in Dairy Cows After Artificial Insemination. Animal Health Prod and Hyg, 2023; 12(2): 36-41.
- Ferguson JD. Nutrition and reproduction in dairy cows. The Veterinary Clinics of North America. Food Animal Practice, 1991; 7: 483-508.
- Forde N., Beltman ME., Duffy GB., Mehta JP., O'Gara P., Roche JF., et al. Changes in the endometrial transcriptome during the bovine estrous cycle: Effect of low circulating progesterone and consequences for conceptus elongation. Biol Reprod, 2011; 84: 266-278.
- Fricke PM., Reynolds LP. and Redmer DA. Effect of human chorionic gonadotropin administered early in the estrous cycle on ovulation and subsequent luteal function in cows. J Anim Sci, 1993; 71: 1242.
- Gaja AO., Hamana K., Kubota C. and Kojima T. Evaluation of the effect of a 3rd GnRH injection administered six days after the 2nd GnRH injection of Ovsynch on the reproductive performance of Japanese black cows. J Vet Sci, 2008; 9: 273-279.
- Geisert RD., Morgan GL., Short Jr EC. and Zavy MT. Endocrine events associated with endometrial function and conceptus development in cattle. Reprod Fertil, 1992; 4: 301-305.
- Geary TW., Ansotegui RP., MacNeil MD., Roberts AJ. and Waterman RC. Effects of flunixin meglumine on pregnancy establishment in beef cattle. J Anim Sci, published online November 20, 2009.
- Guilbault LA., Thatcher WW., Drost M. and Haibel GK. Influence of a physiological infusion of

- prostaglandin  $F2\alpha$  into postpartum cows with partially suppressed endogenous production of prostaglandins: 1. Uterine and ovarian morphological responses. Theriogenology, 1987; 27: 931-946.
- Gumen A. and Wiltbank MC. Length of progesterone exposure needed to resolve large follicle an ovular condition in dairy cows. Theriogenology, 2005; 63: 202-218.
- Hansen PJ., Younquist RS. and Threlfall WR. (eds) Current therapy in large animal theriogenology, 2 nd ed., Saunders-Elsevier, St Louis, MO, 2007; PP: 431-441.
- Hansen PJ. and Arechiga CF. Strategies for managing reproduction in the heat-stressed dairy cow. J Dairy Sci, 1999; 82: 36-50.
- Hansel SW., Larson LL., Laster D., Wagner JF. and Braun RK. Influence of human Chorionic Gonadotropin on Pregnancy rates in lactating dairy and beef cows. J Dairy sci, 1976; 59(4).
- Hosmer DW. and Lemeshow S. Applied logistic regression. New York, USA: Wiley, 1989.
- Inskeep EK. Preovulatory, postovulatory, and post maternal recognition effects of concentrations of progesterone on embryonic survival in the cow. J Anim Sci, 2004; 82 (E-Suppl): E24-E39.
- Kasimanickam R., Gold J., Moore D., Kastelic JP., Pyrdek D. and Ratzgurg K. Injectable or transdermal flunixin meglumine improves pregnancy rates in embryo transfer recipient beef cows without altering to estrus. Theriogenology, 2019; 140: 8-17.
- Kasimanickam R., Hall JB., Estill CT., Kastelic JP., Josheph C., Abdol Aziz RL., et al. flunixin meglumine improves pregnancy rate in embryo recipient beef cows with an excitable temperament. Theriogenology, 2018; 107: 70-77.
- Kamran K., Ekrami ZA., Ghasemzadeh-Nava B., Tajik H., Bolourchi P. and Tamadon MH. Effect of post- insemination of flunixin meglumine on Corpus luteum maintenance and plasma progesterone concentration and pregnancy rate in

- heat stressed Holstein dairy cows. J Anim Vet Adv, 2011; 10 (16): 2176-2180.
- Kidder HE., Black WG., Wiltbank JN., Ulberg LC. and Casida LE. Fertilization rates and embryonic death rates in cows bred to bulls of different levels of fertility. J Dairy Sci, 1954; 691-697.
- Kraevskiy AY., Sokolyuk VM. and Travetskyi MO. Application of Surfagon and Ketaprofen for increasing fertility and preventing embryonic death in cows after insemination. Ukrainian J Ecol, 2020; 10(4): 159-164.
- Lopez H., Satter LD. and Wiltbank MC. Relationship between level of milk production and estrous behavior of lactating dairy cows. Anim Reprod Sci, 2004; 81: 209-223.
- Lucy MC. Reproductive loss in high-producing dairy cattle: Where will it end? J Dairy Sci, 2001; 84: 1277-1293.
- Mann GE., Fray MD. and Lamming GE. Effects of time of progesterone supplementation on embryo development and interferon-tau production in the cow. Vet J, 2006; 171: 500-503.
- Merrill ML., Ansotegui RP., Burns PD., MacNeil MD. and Geary TW. Effects of flunixin meglumine and transportation stress on pregnancy establishment in beef cows. J Anim Sci, 2005; 85: 1547-1554.
- Nishigai M., Kamomae H., Tanaka T. and Kaneda Y. Improvement of pregnancy rate in Japanese Black cows by administration of hCG to recipients of transferred frozen-thawed embryos. Theriogenology, 2002; 58: 1597-1606.
- NRC. Nutrient Requirements of Dairy Cattle, 7th revised edition, Washington, DC: Natl Acad Sci, 2001.
- Odens Vik KH. Pharmacokinetics of flunixin and its effect on prostaglandin F2α metabolite concentrations after oral and intravenous administration in heifers. J Vet Pharmacol Therap, 1995; 18: 254-259.

- Purcell SH., Beal WE. and Gray KR. Effect of a CIDR insert and flunixin meglumine, administered at the time of embryo transfer, on pregnancy rate and resynchronization of estrus in beef cattle. Theriogenology, 2005; 64: 867-878.
- Perry GA., Smith MF., Lucy MC., Green JA., Parks TE., MacNeil MD., et al. Relationship between follicle size at insemination and pregnancy success. Proc Natl Acad Sci, 2005; 102: 5268-5273.
- Raja Mahendran R. and Sianangama PC. Effect of human chorionic gonadotropin on dominant follicles in cows: formation of accessory corpora lutea, progesterone production and pregnancy rates. J Reprod Fertil, 1992; 95:577-584.
- Rensis F., Valentini R., Gorrieri F., Bottarelli E. and Lopez-Gatius F. Inducing ovulation with hCG improves the fertility of dairy cows during the warm season. Theriogenology, 2008; 69(9): 1077-1082.
- Roberts RM., Xie S., and Mathialagan N. Maternal recognition of pregnancy. Biol Reprod, 1996; 54: 294-302.
- Roberts RM., Cross JC. and Leman DW. Interferons as hormones of pregnancy. Endocrine Rev, 1992; 13: 432-452.
- Rossetti RC., Perdigão A., Mesquita FS., Sá Filho M., Nogueira GD.P, Machado R., et al. Effects of flunixin meglumine, recombinant bovine somatotropin and/ or human chorionic gonadotropin on pregnancy rates in Nelore cows. Theriogenology, 2011; 76(4): 751-758.
- Sangsritavong S., Combs DK., Sartori R., Armentano LE. and Wiltbank MC. High feed intake increases liver blood flow and metabolism of progesterone and estradiol-17beta in dairy cattle. J Dairy Sci, 2002; 85: 2831-2842.
- Sianangama PC. and Raja Mahendran R. Effect of human chorionic gonadotropin administered at specific times following breeding on milk progesterone and pregnancy in cows. Theriogenology, 1992; 38:85-96.

- Scenna FN., Hockett ME., Towns TM., Saxton MA., Rohrbach NR., Wehrman ME., et al. Influence of a prostaglandin synthesis inhibitor administered at embryo transfer on pregnancy rates of recipient cows. Prostaglandins Another Lipid Mediat, 2005; 78: 38-45.
- Schmitt EJP., Diaz TM., Barros C., de la Sota RL., Drost M., Fredriksson EW., et al. Differential response of luteal phase and fertility in cattle following ovulation of the first wave follicle with human chorionic gonadotropin or an agonist of gonadotropin- releasing hormone. J Anim Sci, 1996a; 74:1074-1083.
- Schmitt EJ., Barros PCM., Fields PA., Fields MJ., Diaz T., Kluge JM., et al. A cellular and endocrine characterization of the original and induced corpus luteum after administration of a gonadotropin-releasing hormone agonist or human chorionic gonadotropin on day five of the estrous cycle. J Anim Sci., 1996b; 74: 1915-1929.
- Sreenan JM. and Diskin MG. The extent and timing of embryonic mortality in cattle. In: Sree nan JM, Diskin, M.G., editors. Embryonic mortality in farm animals. Brussels: Martinus Nijhoff, 1986; 142-158.
- Satterfield MC., Bazer FW. and Spencer TE. Progesterone regulation of preimplantation conceptus growth and galectin 15 (LGALS15) in the ovine uterus. Biol Reprod, 2006; 75: 289-296.
- Stevenson JS., Phatak AP., Rettmer I. and Stewart RE. Post insemination administration of receptal: Follicular dymics, duration of cycle, hormonal responses, and pregnancy rates. J Dairy Sci, 1993; 76: 2536–2547.
- Stevenson JS., Portaluppi MA., Ten House DE, Lloyd A., Eborn DR., Kacuba S., and et al. Interventions after artificial insemination: conception rates, pregnancy survival, and ovarian responses to gonadotropin-releasing hormone, human chorionic gonadotropin, and progesterone. J Dairy Sci, 2007; 90: 331-340.
- Santos JE., Thatcher WW., Pool L. and Overton MW. Effect of human chorionic gonadotropin on luteal function and reproductive performance of high-

- producing lactating Holstein dairy cows. J Anim Sci, 2001; 79: 2881-2894.
- Shapiro SS. and Wilk M. Analysis of variance test for normality. Biometrika, 1965; 52; 591-601.
- Singh H., Agarwal S., Singh P., Kumar S., Verma PK. and Pandey AK. Effects of hCG administration at day 6 of the estrous cycle on follicular and luteal studies of repeat breeder cross-bred cow. J Entomol Zool Stud, 2020; 8(4): 1326-1330.
- Tahir Karasashin HA., Fatam S., Sukru D. and Huseyn A. effect of flunixin meglumine during and after embryo transfer on pregnancy rate in cattle. 2021, reprod domes anim.
- Thatcher WW., Staples CR., Danet-Desnoyers G., Oldick B. and Schmitt EP. Embryo health and mortality in sheep and cattle. J Anim Sci, 72(3): 1994; 16-30.
- Thatcher WW., Guzeloglu A., Mattos R., Binelli M., Hansen TR. and Pru JK. Uterine conceptus interactions and reproductive failure in cattle. Theriogenology, 2001; 56: 1435-1450.
- Thatcher WW. and Collier RJ. Morrow DA (ed) Current therapy in theriogenology, 2<sup>nd</sup> ed, WB Saunders, Philadelphia, 1986.

- Vasconcelos JL., Sartori R., Oliveira HN., Guenther JG. and Wiltbank MC. Reduction in size of the ovulatory follicle reduces subsequent luteal size and pregnancy rate. Theriogenology, 2001; 56: 307-314.
- Walker RS., Burns PD., Engle TE., Sides GE. and Zalesky DD. Effects of human chorionic gonadotrophin administration on artificial insemination pregnancy rates in beef heifers. Prof Anim Sci, 2005; 21(5): 361-364.
- Wise ME., Rodriguez RE., Armstrong DV., Huber JT., Wiersma F. and Hunter R. Fertility and hormonal responses to temporary relief of heat stress in lactating dairy cows. Theriogenology, 1988; 29: 1027.
- Wiltbank M., Lopez H., Sartori R., Sangsritavong S. and Gumen A. Changes in reproductive physiology of lactating dairy cows due to elevated steroid metabolism. Theriogenology, 2006; 65: 17-29.
- Wolfenson D., Thatcher WW., Badinga L., Savio JD., Meidan R., Lew BJ., et al. Effect of heat stress on follicular development during the estrous cycle in lactating dairy cattle. Biol Reprod, 1995; 52: 1106-1113.

م*قاله پژوهشی* 





# تاثیر گنادوتروپین کوریونی انسانی و فلونکسین مگلومین بر روی میزان آبستنی در گاوهای شیری

نوید اعرابی\*، محمد مهدی هادی پور

گروه علوم بالینی، دانشکده دامپزشکی، واحد کازرون، دانشگاه آزاد اسلامی، کازرون، ایران

تاریخ دریافت: ۱۴۰۲/۰۸/۱۸ اصلاح نهایی: ۱۴۰۲/۰۹/۳۰ تاریخ پذیرش: ۱۴۰۲/۱۰/۰۸

## چکیده

**زمینه و هدف** : افزایش ناباروری و کاهش میزان آبستنی دو مورد از چالش برانگیزترین مسائل در گاوهای شیری مدرن است. هدف از انجام این مطالعه بررسی تاثیر گنادوتروپین کوریونی انسانی روز ۵ و تزریق فلونیکسین مگلومین در روزهای ۱۴–۱۶ بعد از تلقیح بروی آبستنی درگاوهای شیری پرتولید در دو فصل سرد و گرم بود.

مواد و روشها: در این مطالعه تعداد ۵۷۷ راس گاو شیری انتخاب وبه طور تصادفی به چهار گروه در هر دو فصل سرد (تعداد ۳۹۶ گاو) و گرم (تعداد ۱۸۱ گاو) تقسیم شدند. شاخص گرما - رطوبت (۸۵-۹۲) نشان داد که گاو ها در فصل گرم تحت استرس گرمایی بودند. گروه یک: تزریق هورمون گنادوتروپین کوریونی انسانی روز که بعد از تلقیح، گروه دو: تزریق هورمون گنادوتروپین کوریونی انسانی روز پنچم و فلونیکسین مگلومین بین روزهای ۱۹–۱۶ بعد از تلقیح، گروه دون تزریق فلونیکسین روزهای ۵ و ۱۲ بعد از تلقیح نمونه خون اخذ گروه سوم: تزریق فلونیکسین روزهای بین ۱۴–۱۶ و گروه چهارم: گروه کنترل. جهت بررسی میزان پروژسترون در روزهای ۵ و ۱۲ بعد از تلقیح سونو گرافی انجام شد. گردید، همچنین جهت بررسی تعداد جسم زرد، اندازه گیری اقطار جسم زرد در روزهای ۵ و ۱۲ و آبستنی در روز ۲۸–۳۰ بعد از تلقیح سونو گرافی انجام شد. یافته ها: نتایج نشان داد که میزان پروژسترون در گروه های یک و دو بیشتر از دو گروه دیگر در فصل سرد بود ( ۱۳ و ۱۱۲۷، ۱۲۰۱۰) در صورتیکه تفاوت معنی داری در بین گروه ها در فصل گرم مشاهده نشد. همچنین اختلاف معنی داری در بین گروه ها در فصل گرم مشاهده نشد. همچنین اختلاف معنی داری در بین گروه ها در فصل گرم مشاهده نشد. همچنین اختلاف معنی داری در بین گروه ها در فصل گرم از لحاظ آبستنی مشاهده نشد ولی در فصل سرد تغییرات معنی دار بین گروه ها در فصل گرم از لحاظ آبستنی مشاهده نشد ولی در فصل سرد تغییرات معنی دار بین گروه ها در فصل گرم از لحاظ آبستنی مشاهده نشد ولی در فصل سرد تغییرات معنی دار بین گروه ها در فصل گرم از لحاظ آبستنی مشاهده نشد ولی در فصل سرد تغییرات معنی دار بین گروه ها در فصل گرم از لحاظ آبستنی مشاهده نشد ولی در فصل سرد تغییرات معنی دار بین گروه ها در فصل گرم از لحاظ آبستنی مشاهده نشد ولی در فصل سرد تغییرات معنی دار بین گروه ها در فصل گرم از لحاظ آبستنی مشاهده نشد ولی در فصل سرد تغییرات معنی دار بین گروه ها در فصل گرم از لحاظ آبستنی مشاهده نشد ولی در فصل سرد تغییرات معنی دار بین گروه ها در فصل گرم از لحاظ آبستنی مشاهده نشد ولی در فصل سرد بود کروه کرد بین گروه ها در فصل سرد بین گروه ها در فصل سرد بود کرد بین گروه کرد بین گروه ها در فصل سرد بود کرد بین گروه ها در فصل سرد بود کرد

**نتیجه گیری:** با توجه به نتایج حاصل از این مطالعه استفاده از فلونیکسین مگلومین تاثیری در بهبود باروی در دو فصل سرد و گرم نداشت ولی تزریق گنادوتروپین کوریونی انسانی روز ۵ بعد از تلقیح توانست تعداد جسم زرد و میزان پروژسترون و همچنین میزان آبستنی را در فصل سرد افزایش دهد ولی درفصل گرم هیچ تاثیری مشاهده نشد.

واژههای کلیدی: کوریونیک انسانی، فلونیکسین مگلومین، پروژسترون، آبستنی، استرس گرمایی

نوید اعرابی، محمد مهدی هادی پور. تاثیر گنادوتروپین کوریونی انسانی و فلونکسین مگلومین برروی میزان آبستنی در گاوهای شیری. مجله طب دامپزشکی جایگزین. ۱۴۰۳؛ ۷(۲۰)۷: ۱۱۹۴–۱۱۹۴.